

COCHRANE LAKE New Limerick & Smyrna, Aroostook Co. U.S.G.S. Ludlow, Me (7 1/2')

Fishes

Landlocked salmon Brown trout Brook trout Splake White perch Yellow perch Chain pickerel Minnows Golden shiner Redbelly dace Blacknose dace Creek chub Fallfish (chub) White sucker Hornpout (bullhead) Pumpkinseed sunfish American eel

Physical Characteristics

Area - 92 acres Average depth - 13 feet Maximum depth - 33 feet Temperatures: Surface - 75⁰F 33 feet - 42⁰F

Principal fisheries: Splake, white perch, chain pickerel

Cochrane Lake has marginal water quality for coldwater gamefish. Salmon and brown trout management was discontinued due to relatively low returns in the sport fishery. Salmon are likely low in number or nonexistent. Most recently, splake have been stocked to provide a trout fishery. Initially, growth rates of splake were very good since splake are capable of utilizing yellow perch and other fishes as food.

Conservative fishery regulations have been implemented to maintain and enhance sportfishing opportunities.

No public boat launching facility is available at Cochrane Lake, but canoes and small boats can be carried from Route 2 to the outlet for access to the lake.

Surveyed - August 1952 (Revised - 1990, 2000) Maine Department of Inland Fisheries and Wildlife Funded in part by Federal Aid in Fish Restoration Act under Federal Project F-28-P

L1744G