MOOSELOOKMEGUNTIC LAKE Rangeley, Rangeley Plt., Franklin Co. Richardsontown Twp., Adamstown Twp., Oxford County, Maine

Fishes

Salmon M	innows				
Brook trout (squaretail)	Fallfish				
Brown trout	Creek chub				
Smelt	Lake chub				
White sucker	Golden shiner				
Longnose sucker	Common shiner				
Yellow perch	Fathead minnow				
Hornpout (bullhead)	Blacknose dace				
Landlocked alewife	Pearl dace				
	Redbelly dace				

Physical Characteristics

Area - 1	6,300	acres Temperatures					
Maximum	depth	-	139				70°F 45°F

Principal fishery: Salmon, brook trout

Mooselookmeguntic and Cupsuptic Lakes are separate bodies of water that have been joined to make one lake by level of the water raising Mooselookmeguntic Lake by some 14 feet. This water level is maintained by a 20 foot dam at Upper Dam, the outlet of The two basins Mooselookmeguntic Lake. will be considered together in this report and the name Mooselookmeguntic will refer to both lakes.

Mooselookmeguntic Lake, as with all the Rangeley Lakes except Aziscohos, provides excellent temperatures, dissolved oxygen levels, and general water quality characteristics for coldwater fishes. These conditions provide a habitat that produces good salmon and trout fishing despite the presence of yellow perch, a competing warmwater species. Brown trout are rarely taken, as only a relict population exists in the lake.

Mooselookmeguntic Lake has many tributaries which provide excellent spawning and nursery areas for trout and salmon. Kennebago River, Cupsuptic River, and Bemis Stream are the main spawning areas for Mooselookmeguntic Lake and are probably the best spawning streams in the Rangeley area.

Surveyed - July, 1939 (Revised, 1953, 1985, 1992) Maine Department of Inland Fisheries and Wildlife Funded in part by Federal Aid in Fish Restoration Act under Federal Project F-28-P L3302D

