HEALTH IN MAINE: ETHNICITY

This table shows the percentage or rate of each indicator among Hispanics and Non-Hispanics in Maine. For example, in 2015-2019, 18.6% of Hispanics and 11.0% of Non-Hispanics lived in poverty in the state.

The MSCHNA collects and analyzes data on health outcomes, health behaviors, social determinants of health, and demographics. Some subpopulations experiencing health disparities are small, resulting in data that is less reliable due low numbers or unavailable due to privacy concerns. The data that is collected and available is included in these tables. Due to data collection and availability, the list of data in these tables may be shorter than what appears in the County Health Profiles.

The MSCHNA looks forward to supporting the efforts of Maine's Office of Population Health Equity to address these limitations and enhance data collection practices.

POPULATION (MAINE 2019)

HISPANIC 20,714

NON-HISPANIC 1,243,607

Health Indicators by Ethnicity:

| | ETHNICITY | | | |
|--|----------------------|------------------|----------------------|--|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE | |
| DEMOGRAPHICS | | | | |
| Population (percent of total Maine population) | 2019 | 2019 | 2019 | |
| | 1.7% | 98.3% | 100% | |
| Veterans | 2015-2019 | 2015-2019 | 2015-2019 | |
| | 5.8% | 9.8% | 9.6% | |
| Gay, lesbian and bisexual (high school students) | 2019 | 2019 | 2019 | |
| | 15.8% | 12.3% | 12.4% | |
| Gay, lesbian and bisexual (adults) | 2011-2015 | 2011-2015 | 2011-2015 | |
| | & 2017 | & 2017 | & 2017 | |
| | 9.0% | 3.5% | 3.5% | |
| Transgender youth (high school students) | 2019 | 2019 | 2019 | |
| | 5.2% | 1.5% | 1.6% | |
| Persons with a disability | 2015-2019 | 2015-2019 | 2015-2019 | |
| | 13.3% | 16.1% | 16.0% | |
| SOCIAL DETERMINANTS OF HEALTH | | | | |
| Individuals living in poverty | 2015-2019 | 2015-2019 | 2015-2019 | |
| | 18.6% | 11.0% | 11.8% | |
| Median household income | 2015-2019 | 2015-2019 | 2015-2019 | |
| | \$ 49,082 | \$ 58,522 | \$57,918 | |
| High school student graduation | 2020 81.9% | - | 2020 87.4% | |
| Housing insecure (high school students) | 2019 | 2019 | 2019 | |
| | 10.7% | 2.8% | 3.3% | |
| Adverse childhood experiences (high school students) | 2019 | 2019 | 2019 | |
| | 30.9% | 21.0% | 21.3% | |

| | ETHN | IICITY | |
|---|--------------|------------------|--------------|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE |
| GENERAL HEALTH STATUS | | | |
| Fair or poor health (self-rated) | 2011-2017 | 2011-2017 | 2011-2017 |
| | 16.1% | 15.7% | 15.8% |
| 14 or more days lost due to poor physical health | 2011-2017 | 2011-2017 | 2011-2017 |
| | 9.0% | 12.8% | 12.8% |
| 14 or more days lost due to poor mental health | 2011-2017 | 2011-2017 | 2011-2017 |
| | 12.5% | 12.3% | 12.4% |
| Three or more chronic conditions | 2011-2017 | 2011-2017 | 2011-2017 |
| | 9.2% | 14.5% | 14.5% |
| OVERALL MORTALITY | <u>'</u> | | |
| Overall death rate per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 289.3 | 754.5 | 754.7 |
| ACCESS | <u>'</u> | | |
| Uninsured | 2015-2019 | 2015-2019 | 2015-2019 |
| | 13.3% | 7.6% | 7.9% |
| MaineCare enrollment (all ages) | 2020 | 2020 | 2020 |
| | 26.7% | 23.4% | 29.1% |
| MaineCare enrollment (ages 0-19) | 2020 | 2020 | 2020 |
| | 47.0% | 45.5% | 43.8% |
| Usual primary care provider (adults) | 2011-2017 | 2011-2017 | 2011-2017 |
| | 81.9% | 87.9% | 87.8% |
| Primary care visit to any primary care provider in the past year | 2011-2017 | 2011-2017 | 2011-2017 |
| | 66.4% | 71.7% | 71.6% |
| Cost barriers to health care | 2011-2017 | 2011-2017 | 2011-2017 |
| | 19.6% | 10.7% | 10.8% |
| HEALTH CARE QUALITY | | | |
| Ambulatory care-sensitive condition hospitalizations per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 |
| | 34.7 | 59.5 | 61.4 |
| Ambulatory care-sensitive condition emergency department rate per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 |
| | 185.0 | 275.0 | 282.5 |
| CANCER | | | |
| All cancer deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 59.8 | 173.6 | 172.9 |
| Colorectal cancer deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 2.8 | 13.5 | 13.4 |
| Female breast cancer deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 8.9 | 18.2 | 18.1 |
| Lung cancer deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 13.4 | 49.1 | 48.9 |
| Prostate cancer deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 9.2 | 19.5 | 19.5 |
| Tobacco-related cancer deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 18.6 | 53.4 | 53.2 |
| All cancer new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 253.0 | 482.4 | 480.8 |

| | ETHN | IICITY | |
|---|--|--|--------------------------------------|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE |
| CANCER (CONTINUED) | | | |
| Bladder cancer new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 18.0 | 27.5 | 27.4 |
| Colorectal cancer new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 16.9 | 38.2 | 38.0 |
| Colorectal late-stage new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 12.7 | 20.7 | 20.6 |
| Female breast cancer new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 77.6 | 126.8 | 126.4 |
| Female breast cancer late-stage new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 28.3 | 39.6 | 39.5 |
| Lung cancer new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 36.4 | 73.1 | 72.9 |
| Lung cancer late-stage incidence per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 25.7 | 52.9 | 52.7 |
| Melanoma skin cancer new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 5.1 | 25.4 | 25.2 |
| Prostate cancer new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 47.9 | 99.9 | 99.7 |
| Tobacco-related cancer (excluding lung cancer) new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 78.8 | 135.8 | 135.5 |
| HPV-associated cancer new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 5.8 | 13.1 | 13.0 |
| Obesity-associated cancer (excluding colon cancer) new cases per 100,000 population | 2009-2018 | 2009-2018 | 2009-2018 |
| | 86.5 | 133.5 | 133.1 |
| Breast cancer screening up-to-date | 2012, 2014 & 2016 | 2012, 2014 & 2016 82.5% | 2012, 2014 & 2016 82.5% |
| Colorectal cancer screening up-to-date | 2012, 2014 & 2016 | 2012, 2014 & 2016 74.3% | 2012, 2014 & 2016 74.1% |
| Cervical cancer screening up-to-date | 2012, 2014 | 2012, 2014 | 2012, 2014 |
| | & 2016 | & 2016 | & 2016 |
| | 88.9% | 84.9% | 84.9% |
| CARDIOVASCULAR DISEASE | | | |
| High Blood Pressure | 2011, 2013, | 2011, 2013, | 2011, 2013, |
| | 2015 & | 2015 & | 2015 & |
| | 2017 | 2017 | 2017 |
| High cholesterol | 23.8% 2011, 2013, 2015 & 2017 | 33.7% 2011, 2013, 2015 & 2017 | 33.6% 2011, 2013, 2015 & 2017 |
| Cardiovascular disease deaths per 100,000 population | 32.3% | 39.1% | 39.1% |
| | 2010-2019 | 2010-2019 | 2010-2019 |
| | 62.9 | 195.1 | 194.4 |
| Coronary heart disease deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 26.9 | 83.2 | 82.8 |
| Heart attack deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 8.9 | 28.2 | 28.1 |

| | ETHNICITY | | | |
|--|--------------|---------------------------|---------------------------|--|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE | |
| CARDIOVASCULAR DISEASE (CONTINUED) | | | | |
| Stroke deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 | |
| | 17.2 | 34.0 | 33.9 | |
| Heart attack hospitalizations per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 13.6 | 22.2 | 22.5 | |
| Heart failure hospitalizations per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 2.8 | 10.8 | 11.2 | |
| High blood pressure hospitalizations per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 8.5 | 13.7 | 13.8 | |
| Stroke hospitalizations per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 14.8 | 20.7 | 21.2 | |
| Cholesterol checked in past five years | 2011, 2013, | 2011, 2013, | 2011, 2013, | |
| | 2015 & | 2015 & | 2015 & | |
| | 2017 | 2017 | 2017 | |
| DIABETES | 68.4% | 82.3% | 82.1% | |
| Diabetes | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 8.5% | 9.9% | 9.9% | |
| Prediabetes | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 6.7% | 7.6% | 7.6% | |
| Diabetes deaths (underlying cause) per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 | |
| | 18.5 | 21.8 | 21.8 | |
| Diabetes hospitalizations (principal diagnosis) per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 10.7 | 12.4 | 12.7 | |
| Diabetes emergency department rate (principal diagnosis) per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 27.2 | 30.4 | 31.2 | |
| A1c test at least twice/year (adults with diabetes) | 2011-2017 | 2011-2017 77.7% | 2011-2017 77.5% | |
| Formal diabetes education (adults with diabetes) | 2011-2017 | 2011-2017 59.2% | 2011-2017 59.1% | |
| Foot exam annually (adults with diabetes) | 2011-2017 | 2011-2017 83.9% | 2011-2017 83.9% | |
| Dilated eye exam annually (adults with diabetes) | 2011-2017 | 2011-2017 70.4% | 2011-2017 70.4% | |
| RESPIRATORY HEALTH | | | | |
| Current asthma (adults) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 12.7% | 11.6% | 11.6% | |
| Current asthma (youth ages 0-17) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 10.2% | 8.8% | 8.9% | |
| Chronic lower respiratory disease deaths per 100,000 population | 2010-2019 | 2010-2019 48.1 | 2010-2019 48.1 | |
| Asthma emergency department rate per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 38.5 | 41.2 | 42.5 | |
| Chronic obstructive pulmonary disease (COPD) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 2.6% | 8.0% | 7.7% | |
| Chronic obstructive pulmonary disease hospitalizations per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 7.3 | 16.0 | 16.7 | |

| | ETHN | ETHNICITY | | |
|--|--------------|------------------|--------------|--|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE | |
| RESPIRATORY HEALTH (CONTINUED) | | | | |
| Pneumonia hospitalizations per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 9.2 | 18.5 | 19.1 | |
| PHYSICAL ACTIVITY, NUTRITION AND WEIGHT | | | | |
| Obesity (adults) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 28.5% | 28.9% | 28.9% | |
| Overweight (adults) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 34.1% | 36.2% | 36.1% | |
| Obesity (high school students) | 2019 | 2019 | 2019 | |
| | 20.2% | 14.8% | 15.0% | |
| Obesity (middle school students) | 2019 | 2019 | 2019 | |
| | 15.2% | 15.2% | 15.1% | |
| Overweight (high school students) | 2019 | 2019 | 2019 | |
| | 18.6% | 16.2% | 16.3% | |
| Overweight (middle school students) | 2019 | 2019 | 2019 | |
| | 15.6% | 17.5% | 17.3% | |
| Sedentary lifestyle – no leisure-time physical activity in past month (adults) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 20.9% | 22.4% | 22.4% | |
| Met aerobic physical activity recommendations (adults) | 2011, 2013, | 2011, 2013, | 2011, 2013, | |
| | 2015 & | 2015 & | 2015 & | |
| | 2017 | 2017 | 2017 | |
| | 53.8% | 54.3% | 54.2% | |
| Met physical activity recommendations (high school students) | 2019 | 2019 | 2019 | |
| | 21.5% | 20.9% | 20.9% | |
| Met physical activity recommendations (middle school students) | 2019 | 2019 | 2019 | |
| | 25.1% | 25.5% | 25.5% | |
| Fewer than two hours combined screen time (high school students) | 2019 | 2019 | 2019 | |
| | 38.7% | 33.6% | 33.9% | |
| Fewer than two hours combined screen time (middle school students) | 2019 | 2019 | 2019 | |
| | 26.3% | 28.0% | 28.2% | |
| Fruit consumption (adults reporting less than one serving per day) | 2017 | 2017 | 2017 | |
| | - | 31.4% | 31.4% | |
| Vegetable consumption (adults reporting less than one serving per day) | 2017 | 2017 | 2017 | |
| | - | 12.4% | 12.4% | |
| Fruit and vegetable consumption (high school students reporting 5 or more a day) | 2019 | 2019 | 2019 | |
| | 17.9% | 15.1% | 15.2% | |
| Fruit and vegetable consumption (middle school students reporting 5 or more a day) | 2019 | 2019 | 2019 | |
| | 22.0% | 20.9% | 20.9% | |
| Soda/sports drink consumption (high school students reporting 1 or more a day) | 2019 | 2019 | 2019 | |
| | 22.0% | 19.4% | 19.6% | |
| Soda/sports drink consumption (middle school students reporting 1 or more a day) | 2019 | 2019 | 2019 | |
| | 19.6% | 17.5% | 17.7% | |

| Infant deaths per 1,000 live births 2015-2019 5.8 5.8 5.8 Low birth weight (<2500 grams) 2017-2019 | | EIRIN | ETHNICITY | |
|---|---|-----------|-----------|-------|
| Infant deaths per 1,000 live births 5.9 5.8 5.9 5.8 5.8 5.8 5.8 5.8 5.8 5.8 5.8 5.9 5.8 5.8 5.8 5.8 5.9 5.8 5.8 5.8 5.8 5.9 5.8 5.8 5.8 5.9 5.8 5.8 5.8 5.9 5.8 5.9 5.8 5.8 5.9 5.9 5.8 5.8 5.8 5.9 5.9 5.8 5.8 5.8 5.9 5.9 5.8 5.9 5.8 5.9 5.9 5.8 5.9 5.8 5.9 5.8 5.9 5.8 5.9 5.9 5.8 5.9 5.9 5.8 5.8 5.8 5.9 5.9 5.8 5.9 5.8 5.8 5.8 5.8 5.8 5.9 5.9 5.8 5.9 5.9 5.8 5.9 5.9 5.8 5.9 5.9 5.8 5.9 5.9 5.8 5.9 5.9 5.9 5.9 5.9 5.8 5.9 5.9 5.9 5.9 5.8 5.9 5.9 5.9 5.8 5.9 5.9 5.9 5.8 5.9 5.9 5.9 5.8 5.9 5.9 5.9 5.8 5.9 5.9 5.8 5.9 5. | INDICATOR | HISPANIC | - | MAINE |
| Infant deaths per 1,000 live births 5,9 5,8 5,8 Low birth weight (<2500 grams) | PREGNANCY AND BIRTH OUTCOMES | | | |
| Low birth weight (<2500 grams) 7.9% 7.2% 7.3% 7.3% 2017-2019 2017-2019 2018-2019 2017-2019 2018-2019 201 | Infant deaths per 1,000 live births | | | |
| Pre-term live births 7.5% 8.8% 2015-2019 | Low birth weight (<2500 grams) | | | |
| Births to 15-19 year olds per 1,000 population 14.1 12.5 10.0 10.5 2015-2019 | Pre-term live births | | | |
| Dinitended births 24.3% 21.5% 21.6% | Births to 15-19 year olds per 1,000 population | | | |
| prenatal visits 75.9% 81.9% 82.7% Smoked during pregnancy 2017-2019 2017-2019 2018-2019 Drank alcohol during pregnancy 2015-2019 2015-2019 2015-2019 Infants who are ever breast fed 2017-2019 2017-2019 2018-2019 C-sections among low-risk first births 2017-2017 2017-2019 2017-2019 2018-2019 C-sections among low-risk first births 2017-2017 2017-2019 2010 2010 C-sections among low-risk first births 2012 2020 2020 2020 C-sections among low-risk first births 2012 2017 2017-2017 2017-2017 2017-2017 </td <td>Unintended births</td> <td></td> <td></td> <td></td> | Unintended births | | | |
| Smoked during pregnancy 8.9% 12.4% 11.9% Drank alcohol during pregnancy 2015-2019 2015-2019 2015-2019 2015-2019 2015-2019 2015-2019 2015-2019 2015-2019 2017-2019 2018-2019 2011-2017 </td <td>•</td> <td></td> <td></td> <td></td> | • | | | |
| Drank alcohol during pregnancy 9.7% 8.5% 8.5% 8.5% 1 | Smoked during pregnancy | | | |
| Infants who are ever breast fed 88.8% 88.4% 89.3% 2017-2019 2017-2019 2018-2 | Drank alcohol during pregnancy | | | |
| C-sections among low-risk first births 26.5% 24.7% 25.2% | Infants who are ever breast fed | | | |
| Developmental screening for MaineCare members 13.2% 16.1% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.6% 16.1% 16.6% 16.1% 16.6% 16.1% 16.6% 16.1% 16.6% 16.1 | C-sections among low-risk first births | | | |
| Table Tabl | CHILDREN WITH SPECIAL HEALTH CARE NEEDS | · | | |
| Arthritis 24.5% 32.2% 31.0% 24.5% 32.2% 31.0% 32.2% 31.0% 2015 & 2015 & 2015 & 2015 & 2015 & 2015 & 2017 & | Developmental screening for MaineCare members | | | |
| Arthritis 24.5% 32.2% 31.0% | COGNITIVE HEALTH | · | | |
| Caregiving at least 20 hours per week 2017 8.9% 4.7% 4.8% | Arthritis | | | |
| Influenza vaccination in the past year (adults) 2011-2017 31.6% 42.4% 42.3% 42.3% 42.4% 42.3% 42.3% 2011-2017 2011-2017 75.2% | Caregiving at least 20 hours per week | 2017 | 2017 | 2017 |
| Name | IMMUNIZATIONS | | 1 | |
| Pineumococcal pneumonia vaccination (adults ages 65+) - 75.2% 75.2% | Influenza vaccination in the past year (adults) | | | |
| Gastrointestinal disease new cases per 100,000 population 2020 2020 2020 2020 2020 2020 2020 202 | Pneumococcal pneumonia vaccination (adults ages 65+) | 2011-2017 | | |
| Hepatitis B (acute) new cases per 100,000 population 16.9 38.3 33.5 Hepatitis A (acute) new cases per 100,000 population 2020 2020 2020 Hepatitis B (acute) new cases per 100,000 population 2020 2020 2020 Hepatitis B (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (acute) new cases per 100,000 population 2020 2020 2020 Hepatitis C (acute) new cases per 100,000 population 25.3 12.5 15.3 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 2020 Hepatitis C (chronic) new cases per 100,000 population 2020 2 | INFECTIOUS DISEASE | | | |
| Hepatitis A (acute) new cases per 100,000 population 0.0 10.5 10.8 | Gastrointestinal disease new cases per 100,000 population | | | |
| Hepatitis B (acute) new cases per 100,000 population 0.0 2.3 3.0 | Hepatitis A (acute) new cases per 100,000 population | | | |
| Hepatitis B (chronic) new cases per 100,000 population 0.0 6.7 9.2 Hepatitis C (acute) new cases per 100,000 population 2020 25.3 2020 12.5 2020 15.3 Hepatitis C (chronic) new cases per 100,000 population 2020 2020 2020 2020 | Hepatitis B (acute) new cases per 100,000 population | | | |
| Hepatitis C (acute) new cases per 100,000 population 25.3 12.5 15.3 Hepatitis C (chronic) new cases per 100,000 population | Hepatitis B (chronic) new cases per 100,000 population | | | |
| Henatitis C (chronic) new cases per 100 000 population | Hepatitis C (acute) new cases per 100,000 population | | | |
| | Hepatitis C (chronic) new cases per 100,000 population | | | |

ETHNICITY

| | ETHN | IICITY | |
|---|-------------------------------------|---|---|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE |
| INFECTIOUS DISEASE (CONTINUED) | | | |
| Lyme disease new cases per 100,000 population | 2020 | 2020 | 2020 |
| | 12.7 | 36.9 | 83.8 |
| Pertussis new cases per 100,000 population | 2020 | 2020 | 2020 |
| | 4.2 | 1.4 | 2.2 |
| Tuberculosis new cases per 100,000 population | 2020 | 2020 | 2020 |
| | 4.2 | 1.2 | 1.3 |
| Chlamydia new cases per 100,000 population | 2020 | 2020 | 2020 |
| | 206.8 | 134.3 | 257.8 |
| Gonorrhea new cases per 100,000 population | 2020 | 2020 | 2020 |
| | 63.3 | 36.6 | 38.7 |
| HIV new cases per 100,000 population | 2020 | 2020 1.2 | 2020 1.2 |
| Syphilis new cases per 100,000 population | 2020 | 2020 | 2020 |
| | 21.1 | 4.5 | 4.9 |
| UNINTENTIONAL INJURY | | | |
| Injury deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 34.1 | 72.2 | 71.7 |
| Fall-related deaths (unintentional) per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 2.8 | 11.1 | 11.0 |
| Poisoning deaths (unintentional and undetermined intent) per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 13.2 | 20.0 | 19.9 |
| Motor vehicle traffic crash (unintentional) deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 5.5 | 11.0 | 10.9 |
| Fall-related injury (unintentional) emergency department rate per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 |
| | 168.3 | 299.8 | 307.4 |
| Traumatic brain injury emergency department rate per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 |
| | 22.6 | 38.2 | 39.2 |
| Always wear seatbelt (high school students) | 2019 | 2019 | 2019 |
| | 59.2% | 70.4% | 70.0% |
| Always wear seatbelt (middle school students) | 2019 | 2019 | 2019 |
| | 65.1% | 78.1% | 77.9% |
| INTENTIONAL INJURY | | | |
| Firearm deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 5.3 | 9.9 | 9.8 |
| Suicide deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 |
| | 6.8 | 16.8 | 16.6 |
| Rape/non-consensual sex (among females, lifetime) | 2011, 2012, 2014, 2016 & 2017 | 2011, 2012, 2014, 2016 & 2017 14.8% | 2011, 2012, 2014, 2016 & 2017 14.9% |
| Violence by current or former intimate partners in past 12 months (among females) | 2011, 2012, 2014 & 2016 | 2011, 2012, 2014 & 2016 1.5% | 2011, 2012, 2014 & 2016 1.5% |
| | 2019 | 2019 | 2019 |

| | ETHNICITY | | | |
|--|--------------|-------------------------|-------------------------|--|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE | |
| INTENTIONAL INJURY (CONTINUED) | | | | |
| Bullied on school property (middle school students) | 2019 | 2019 | 2019 | |
| | 54.2% | 45.9% | 46.3% | |
| Intentional self-injury (high school students) | 2019 | 2019 | 2019 | |
| | 25.3% | 18.4% | 18.7% | |
| Intentional self-injury (middle school students) | 2019 | 2019 | 2019 | |
| | 31.7% | 18.7% | 18.9% | |
| MENTAL HEALTH | | | | |
| Mental health emergency department rate per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 105.9 | 178.5 | 181.5 | |
| Depression, current symptoms (adults) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 16.0% | 9.5% | 9.6% | |
| Depression, lifetime | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 28.5% | 23.5% | 23.6% | |
| Anxiety, lifetime | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 21.8% | 20.2% | 20.3% | |
| Sad/hopeless for two weeks in a row (high school students) | 2019 | 2019 | 2019 | |
| | 39.5% | 31.9% | 32.1% | |
| Sad/hopeless for two weeks in a row (middle school students) | 2019 | 2019 | 2019 | |
| | 39.2% | 24.6% | 24.8% | |
| Seriously considered suicide (high school students) | 2019 | 2019 | 2019 | |
| | 22.3% | 16.2% | 16.4% | |
| Seriously considered suicide (middle school students) | 2019 | 2019 | 2019 | |
| | 30.0% | 19.8% | 19.8% | |
| Chronic disease among persons with depression | 2011-2017 | 2011-2017 | 2011-2017 | |
| | - | 30.9% | 30.8% | |
| Currently receiving outpatient mental health treatment (adults) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 19.5% | 17.8% | 17.8% | |
| ORAL HEALTH | | | | |
| Dentist visits in the past year (adults) | 2012, 2014 | 2012, 2014 | 2012, 2014 | |
| | & 2016 | & 2016 | & 2016 | |
| | 62.7% | 64.3% | 64.2% | |
| Adult tooth loss | 2012, 2014 | 2012, 2014 | 2012, 2014 | |
| | & 2016 | & 2016 | & 2016 | |
| | 11.1% | 19.8% | 19.8% | |
| Ambulatory care sensitive dental emergency department rates for adults per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 80.4 | 134.7 | 136.9 | |
| Ambulatory care sensitive dental emergency department rates for children per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 13.0 | 17.2 | 17.9 | |
| SUBSTANCE USE | | | | |
| Overdose deaths per 100,000 population | 2020 | 2020 | 2020 | |
| | 31.7 | 37.8 | 37.3 | |
| Drug-induced deaths per 100,000 population | 2010-2019 | 2010-2019 | 2010-2019 | |
| | 12.6 | 21.4 | 21.4 | |
| Alcohol-induced deaths per 100,000 population | 2010-2019 | 2010-2019 9.8 | 2010-2019 9.8 | |

ETHNICITY

| | ETHN | IICITY | | |
|---|--------------|------------------|--------------|--|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE | |
| SUBSTANCE USE (CONTINUED) | | THIST AIRIC | | |
| Chronic heavy drinking (adults) | 2013-2017 | 2013-2017 | 2013-2017 | |
| | 9.8% | 7.9% | 8.0% | |
| Binge drinking (adults) | 2011-2017 | 2011-2017 | 2011-2017 | |
| | 21.5% | 17.5% | 17.6% | |
| Past-30-day marijuana use (adults) | 2013-2017 | 2013-2017 | 2013-2017 | |
| | 18.7% | 11.5% | 11.6% | |
| Past-30-day misuse of prescription drugs (adult) | 2013-2017 | 2013-2017 | 2013-2017 | |
| | 1.1% | 1.0% | 1.0% | |
| Past-30-day alcohol use (high school students) | 2019 | 2019 | 2019 | |
| | 28.5% | 22.7% | 22.9% | |
| Past-30-day alcohol use (middle school students) | 2019 | 2019 | 2019 | |
| | 8.1% | 4.0% | 4.0% | |
| Binge drinking (high school students) | 2015 | 2015 | 2015 | |
| | 12.4% | 12.8% | 12.2% | |
| Binge drinking (middle school students) | 2019 | 2019 | 2019 | |
| | 3.2% | 1.2% | 1.3% | |
| Past-30-day marijuana use (high school students) | 2019 | 2019 | 2019 | |
| | 29.3% | 21.8% | 22.1% | |
| Past-30-day marijuana use (middle school students) | 2019 | 2019 | 2019 | |
| | 8.0% | 4.0% | 4.1% | |
| Past-30-day misuse of prescription drugs (high school students) | 2019 | 2019 | 2019 | |
| | 11.9% | 4.6% | 5.0% | |
| Past-30-day misuse of prescription drugs (middle school students) | 2019 | 2019 | 2019 | |
| | 7.3% | 2.8% | 3.0% | |
| Opiate poisoning emergency department rate per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 4.6 | 9.8 | 9.9 | |
| Opiate poisoning hospitalizations per 10,000 population | 2016-2018 | 2016-2018 | 2016-2018 | |
| | 0.4 | 1.4 | 1.4 | |
| TOBACCO USE | | | | |
| Current (every day or somedays) smoking (adults) | 2013-2017 | 2013-2017 | 2013-2017 | |
| | 25.2% | 19.1% | 19.2% | |
| Current (every day or somedays) e-cigarette use (adults) | 2015-2017 | 2015-2017 | 2015-2017 | |
| | 8.8% | 3.4% | 3.5% | |
| Past-30-day cigarette smoking (high school students) | 2019 | 2019 | 2019 | |
| | 14.1% | 6.7% | 7.1% | |
| Past-30-day cigarette smoking (middle school students) | 2019 | 2019 | 2019 | |
| | 3.5% | 1.3% | 1.5% | |
| Past-30-day e-cigarette use (high school students) | 2019 | 2019 | 2019 | |
| | 33.4% | 28.5% | 28.7% | |
| Past-30-day e-cigarette use (middle school students) | 2019 | 2019 | 2019 | |
| | 12.8% | 7.0% | 7.0% | |
| Past-30-day tobacco use (high school students) | 2019 | 2019 | 2019 | |
| | 18.3% | 10.1% | 10.6% | |
| Past-30-day tobacco use (middle school students) | 2019 | 2019 | 2019 | |
| | 5.9% | 3.0% | 3.1% | |
| Environmental tobacco smoke exposure (high school students) | 2019 | 2019 | 2019 | |
| | 32.9% | 26.7% | 27.0% | |
| | 32.3/0 | 20.7/0 | 27.0/0 | |

| | ETHNICITY | | |
|---|----------------------|----------------------|----------------------|
| INDICATOR | HISPANIC | NON- HISPANIC | MAINE |
| TOBACCO USE (CONTINUED) | | | |
| Environmental tobacco smoke exposure (middle school students) | 2019 27.7% | 2019 21.9% | 2019 22.1% |
| Maine QuitLink users | 2020 2.1% | 2020 1.5% | 2020 1.8% |

The Maine Shared Community Health Needs Assessment (Maine Shared CHNA) is a collaboration between Central Maine Healthcare (CMHC), Maine Center for Disease Control and Prevention (Maine CDC), MaineGeneral Health (MGH), MaineHealth (MH), and Northern Light Health (NLH).

The vision of the Maine Shared CHNA is to turn health data into action so that Maine will become the healthiest state in the U.S. The mission of the Maine Shared CHNA is to:

- · Create Shared CHNA Reports,
- Engage and activate communities, and
- Support data-driven health improvements for Maine people.

For more information on Maine Shared CHNA, including a list of data sources and definitions, please visit our web page: www.mainechna.org.









