





Errata sheet

Maine Leading Causes of Death report 2018

March 31, 2022



| Table 6. Leading Causes of Death by County, Maine, 2018 (page 1 of 5) | | | | |
|---|------------------|------------|-------------------|---------------|
| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
| | | | Rate | 95% CI |
| Androscoggin | | | | |
| 1 Cancer (C00-C97) | 258 | 239.6 | 178.5 | 156.3 - 200.7 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 231 | 214.5 | 163.4 | 142.1 - 184.7 |
| 3 Alzheimer's disease (G30) | 101 | 93.8 | 70.4 | 56.7 - 84.2 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 80 | 74.3 | 66.3 | 52.1 - 83.3 |
| 5 Chronic lower respiratory disease (J40-J47) | 75 | 85.4 | 64.7 | 52.1 - 79.6 |
| 6 Cerebrovascular disease (I60-I69) | 46 | 42.7 | 31.3 | 22.9 - 41.9 |
| 7 Diabetes (E10-E14) | 36 | 33.4 | 25.7 | 17.8 - 35.9 |
| 8 Influenza and pneumonia (J10-J18) | 31 | 28.8 | 20.9 | 14.1 - 29.8 |
| 9 Chronic liver disease and cirrhosis (K70,K73-K74) | 24 | 22.3 | 16.9 | 10.6 - 25.6 |
| 10 Suicide (X60-X84,Y87.0) | 19 | DNR | DNR | 16.0 - 20.7 |
| Aroostook | | | | |
| 1 Cancer (C00-C97) | 177 | 263.7 | 156.2 | 131.9 - 180.6 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 169 | 251.8 | 138.9 | 117.4 - 160.4 |
| 3 Chronic lower respiratory disease (J40-J47) | 76 | 113.2 | 62.9 | 49.2 - 79.2 |
| 4 Cerebrovascular disease (I60-I69) | 47 | 70.0 | 39.5 | 28.98 - 52.7 |
| 5 Unintentional injury (V01-X59,Y85-Y86) | 39 | 58.1 | 43.6 | 29.6 - 61.8 |
| 6 Alzheimer's disease (G30) | 30 | 44.7 | 23.2 | 15.6 - 33.1 |
| 7 Diabetes (E10-E14) | 28 | 41.7 | 22.9 | 15.1 - 33.4 |
| 8 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 27 | 40.2 | 20.8 | 13.7 - 30.3 |
| 9 Influenza and pneumonia (J10-J18) | 15 | 22.4 | 12.0 | 6.7 - 19.8 |
| 10 Chronic liver disease and cirrhosis (K70,K73-K74) | 15 | DNR | DNR | 7.8 - 25.0 |
| Cumberland | | | | |
| 1 Cancer (C00-C97) | 574 | 195.5 | 143.2 | 131.3 - 155.1 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 560 | 190.8 | 135.5 | 124.1 - 146.8 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 178 | 60.6 | 52.5 | 44.5 - 60.5 |
| 4 Chronic lower respiratory disease (J40-J47) | 148 | 50.4 | 35.2 | 29.5 - 41.0 |
| 5 Cerebrovascular disease (I60-I69) | 131 | 44.6 | 30.8 | 25.4 - 36.1 |
| 6 Alzheimer's disease (G30) | 106 | 36.1 | 25.3 | 20.5 - 30.1 |
| 7 Diabetes (E10-E14) | 83 | 28.3 | 19.5 | 15.5 - 24.3 |
| 8 Influenza and pneumonia (J10-J18) | 56 | 19.1 | 13.3 | 10.0 - 17.3 |
| 9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 55 | 18.7 | 12.7 | 9.6 - 16.6 |
| 10 Parkinson's disease (G20-G21) | 52 | 17.7 | 12.4 | 9.2 - 16.3 |

| Table 6. Leading Causes of Death by County, Maine, 2018 (page 1 of 5) | | | | |
|---|------------------|------------|-------------------|---------------|
| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
| | | | Rate | 95% CI |
| Androscoggin | | | | |
| 1 Cancer (C00-C97) | 258 | 239.6 | 178.5 | 156.3 - 200.7 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 231 | 214.5 | 163.4 | 142.1 - 184.7 |
| 3 Alzheimer's disease (G30) | 101 | 93.8 | 70.4 | 56.7 - 84.2 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 80 | 74.3 | 66.3 | 52.1 - 83.3 |
| 5 Chronic lower respiratory disease (J40-J47) | 75 | 85.4 | 64.7 | 52.1 - 79.6 |
| 6 Cerebrovascular disease (I60-I69) | 46 | 42.7 | 31.3 | 22.9 - 41.9 |
| 7 Diabetes (E10-E14) | 36 | 33.4 | 25.7 | 17.8 - 35.9 |
| 8 Influenza and pneumonia (J10-J18) | 31 | 28.8 | 20.9 | 14.1 - 29.8 |
| 9 Chronic liver disease and cirrhosis (K70,K73-K74) | 24 | 22.3 | 16.9 | 10.6 - 25.6 |
| 10 Suicide (X60-X84,Y87.0) | 19 | DNR | DNR | 16.0 - 20.7 |
| Aroostook | | | | |
| 1 Cancer (C00-C97) | 177 | 263.7 | 156.2 | 131.9 - 180.6 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 169 | 251.8 | 138.9 | 117.4 - 160.4 |
| 3 Chronic lower respiratory disease (J40-J47) | 76 | 113.2 | 62.9 | 49.2 - 79.2 |
| 4 Cerebrovascular disease (I60-I69) | 47 | 70.0 | 39.5 | 28.98 - 52.7 |
| 5 Unintentional injury (V01-X59,Y85-Y86) | 39 | 58.1 | 43.6 | 29.6 - 61.8 |
| 6 Alzheimer's disease (G30) | 30 | 44.7 | 23.2 | 15.6 - 33.1 |
| 7 Diabetes (E10-E14) | 28 | 41.7 | 22.9 | 15.1 - 33.4 |
| 8 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 27 | 40.2 | 20.8 | 13.7 - 30.3 |
| 9 Influenza and pneumonia (J10-J18) | 15 | 22.4 | 12.0 | 6.7 - 19.8 |
| 10 Chronic liver disease and cirrhosis (K70,K73-K74) | 15 | DNR | DNR | 7.8 - 25.0 |
| Cumberland | | | | |
| 1 Cancer (C00-C97) | 574 | 195.5 | 143.2 | 131.3 - 155.1 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 560 | 190.8 | 135.5 | 124.1 - 146.8 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 178 | 60.6 | 52.5 | 44.5 - 60.5 |
| 4 Chronic lower respiratory disease (J40-J47) | 148 | 50.4 | 35.2 | 29.5 - 41.0 |
| 5 Cerebrovascular disease (I60-I69) | 131 | 44.6 | 30.8 | 25.4 - 36.1 |
| 6 Alzheimer's disease (G30) | 106 | 36.1 | 25.3 | 20.5 - 30.1 |
| 7 Diabetes (E10-E14) | 83 | 28.3 | 19.5 | 15.5 - 24.3 |
| 8 Influenza and pneumonia (J10-J18) | 56 | 19.1 | 13.3 | 10.0 - 17.3 |
| 9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 55 | 18.7 | 12.7 | 9.6 - 16.6 |
| 10 Parkinson's disease (G20-G21) | 52 | 17.7 | 12.4 | 9.2 - 16.3 |

| Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 2 of 5) | | | | |
|---|------------------|------------|-------------------|---------------|
| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
| | | | Rate | 95% CI |
| Franklin | | | | |
| 1 Cancer (C00-C97) | 96 | 321.1 | 200.5 | 161.5 - 246.2 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 58 | 194.0 | 125.1 | 94.5 - 162.5 |
| 3 Chronic lower respiratory disease (J40-J47) | 25 | 83.6 | 44.3 | 27.4 - 68.7 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 23 | 76.9 | 68.6 | 41.3 - 107.2 |
| 5 Cerebrovascular disease (I60-I69) | 20 | DNR | DNR | 24.4 - 61.6 |
| 6 Diabetes (E10-E14) | 15 | DNR | DNR | 17.4 - 53.5 |
| 7 Alzheimer's disease (G30) | 15 | DNR | DNR | 18.3 - 53.9 |
| 8 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 13 | DNR | DNR | 15.1 - 48.6 |
| 9 Influenza and pneumonia (J10-J18) | 11 | DNR | DNR | 12.1 - 50.2 |
| Hancock | | | | |
| 1 Cancer (C00-C97) | 142 | 259.1 | 157.0 | 130.0 - 184.0 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 108 | 197.0 | 117.2 | 94.3 - 140.1 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 40 | 73.0 | 57.9 | 39.6 - 81.6 |
| 4 Chronic lower respiratory disease (J40-J47) | 39 | 71.2 | 40.9 | 28.9 - 56.1 |
| 5 Cerebrovascular disease (I60-I69) | 27 | 49.3 | 26.3 | 17.3 - 38.3 |
| 6 Alzheimer's disease (G30) | 18 | DNR | DNR | 11.0 - 29.4 |
| 7 Diabetes (E10-E14) | 17 | DNR | DNR | 11.9 - 35.0 |
| 8 Parkinson's disease (G20-G21) | 14 | DNR | DNR | 7.9 - 24.2 |
| 9 Influenza and pneumonia (J10-J18) | 12 | DNR | DNR | 5.9 - 19.8 |
| 10 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 10 | DNR | DNR | 4.6 - 17.8 |
| Kennebec | | | | |
| 1 Cancer (C00-C97) | 273 | 223.6 | 150.9 | 132.5 - 169.2 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 273 | 223.6 | 152.3 | 133.9 - 170.7 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 102 | 83.5 | 75.8 | 61.0 - 93.1 |
| 4 Cerebrovascular disease (I60-I69) | 88 | 72.9 | 49.8 | 39.7 - 61.6 |
| 5 Chronic lower respiratory disease (J40-J47) | 89 | 50.8 | 34.1 | 26.0 - 43.9 |
| 6 Diabetes (E10-E14) | 48 | 39.3 | 26.2 | 19.3 - 34.7 |
| 7 Alzheimer's disease (G30) | 24 | 19.7 | 20.0 | 12.5 - 30.3 |
| 8 Suicide (X60-X84,Y87.0) | 24 | 19.7 | 13.0 | 8.3 - 19.3 |
| 9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24 | 19.7 | 13.0 | 8.3 - 19.3 |
| 10 Influenza and pneumonia (J10-J18) | 23 | 18.8 | 12.4 | 7.8 - 18.5 |

| Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 2 of 5) | | | | |
|---|------------------|------------|-------------------|---------------|
| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
| | | | Rate | 95% CI |
| Franklin | | | | |
| 1 Cancer (C00-C97) | 96 | 321.1 | 200.5 | 161.5 - 246.2 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 58 | 194.0 | 125.1 | 94.5 - 162.5 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 23 | 76.9 | 68.5 | 41.2 - 106.9 |
| 4 Chronic lower respiratory disease (J40-J47) | 22 | 73.6 | 44.3 | 27.4 - 67.8 |
| 5 Cerebrovascular disease (I60-I69) | 20 | DNR | DNR | 24.4 - 61.6 |
| 6 Alzheimer's disease (G30) | 15 | DNR | DNR | 18.3 - 53.9 |
| 7 Diabetes (E10-E14) | 15 | DNR | DNR | 17.4 - 53.5 |
| 8 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 13 | DNR | DNR | 15.1 - 48.6 |
| 9 Influenza and pneumonia (J10-J18) | 11 | DNR | DNR | 12.0 - 49.8 |
| Hancock | | | | |
| 1 Cancer (C00-C97) | 142 | 259.1 | 156.9 | 129.9 - 183.9 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 108 | 197.0 | 117.2 | 94.3 - 140.1 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 40 | 73.0 | 57.8 | 39.6 - 81.6 |
| 4 Chronic lower respiratory disease (J40-J47) | 39 | 71.2 | 40.9 | 28.9 - 56.1 |
| 5 Cerebrovascular disease (I60-I69) | 27 | 49.3 | 26.3 | 17.3 - 38.3 |
| 6 Alzheimer's disease (G30) | 18 | DNR | DNR | 11.0 - 29.4 |
| 7 Diabetes (E10-E14) | 17 | DNR | DNR | 11.9 - 35.0 |
| 8 Parkinson's disease (G20-G21) | 14 | DNR | DNR | 7.9 - 24.2 |
| 9 Influenza and pneumonia (J10-J18) | 12 | DNR | DNR | 5.9 - 19.8 |
| 10 Suicide (X60-X84,Y87.0) | 10 | DNR | DNR | 6.3 - 26.3 |
| Kennebec | | | | |
| 1 Heart disease (I00-I09,I11,I13,I20-I51) | 273 | 223.6 | 152.3 | 133.9 - 170.7 |
| 2 Cancer (C00-C97) | 273 | 223.6 | 150.9 | 132.5 - 169.2 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 102 | 83.5 | 75.8 | 61.0 - 93.1 |
| 4 Chronic lower respiratory disease (J40-J47) | 89 | 72.9 | 49.8 | 39.7 - 61.6 |
| 5 Cerebrovascular disease (I60-I69) | 88 | 72.1 | 49.8 | 39.7 - 61.6 |
| 6 Diabetes (E10-E14) | 48 | 39.3 | 26.2 | 19.3 - 34.7 |
| 7 Alzheimer's disease (G30) | 24 | 19.7 | 20.0 | 12.5 - 30.3 |
| 8 Suicide (X60-X84,Y87.0) | 24 | 19.7 | 13.0 | 8.3 - 19.3 |
| 9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24 | 19.7 | 13.0 | 8.3 - 19.3 |
| 10 Influenza and pneumonia (J10-J18) | 23 | 18.8 | 12.4 | 7.8 - 18.5 |

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 3 of 5)

| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
|--|------------------|------------|-------------------|---------------|
| | | | Rate | 95% CI |
| Knox | | | | |
| 1 Cancer (C00-C97) | 134 | 336.9 | 179.9 | 148.9 - 210.9 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 112 | 281.6 | 154.5 | 125.5 - 183.4 |
| 3 Chronic lower respiratory disease (J40-J47) | 32 | 80.5 | 44.5 | 29.8 - 63.9 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 27 | 67.9 | 50.6 | 32.1 - 76.0 |
| 5 Alzheimer's disease (G30) | 21 | 52.8 | 28.7 | 17.8 - 43.9 |
| 6 Cerebrovascular disease (I60-I69) | 21 | 52.8 | 28.4 | 17.3 - 43.9 |
| 7 Diabetes (E10-E14) | 10 | DNR | DNR | 6.9 - 28.7 |
| Lincoln | | | | |
| 1 Heart disease (I00-I09,I11,I13,I20-I51) | 98 | 285.4 | 153.0 | 122.5 - 188.7 |
| 2 Cancer (C00-C97) | 96 | 279.5 | 138.4 | 111.3 - 170.1 |
| 3 Cerebrovascular disease (I60-I69) | 31 | 90.3 | 44.9 | 30.5 - 63.7 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 22 | 64.1 | 54.3 | 31.0 - 88.2 |
| 5 Chronic lower respiratory disease (J40-J47) | 20 | 58.2 | 29.4 | 18.0 - 45.5 |
| 6 Alzheimer's disease (G30) | 20 | DNR | DNR | 18.0 - 45.5 |
| 7 Diabetes (E10-E14) | 11 | DNR | DNR | 7.9 - 28.4 |
| Oxford | | | | |
| 1 Cancer (C00-C97) | 162 | 281.2 | 180.4 | 151.4 - 209.4 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 118 | 204.8 | 135.5 | 110.1 - 160.8 |
| 3 Chronic lower respiratory disease (J40-J47) | 57 | 98.9 | 62.4 | 47.0 - 81.2 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 50 | 86.8 | 80.4 | 58.2 - 108.3 |
| 5 Diabetes (E10-E14) | 30 | 52.1 | 32.6 | 21.9 - 46.9 |
| 6 Cerebrovascular disease (I60-I69) | 23 | 39.9 | 27.2 | 17.1 - 41.3 |
| 7 Alzheimer's disease (G30) | 21 | 36.4 | 24.3 | 15.1 - 37.2 |
| 8 Influenza and pneumonia (J10-J18) | 18 | DNR | DNR | 11.3 - 31.0 |
| 9 Suicide (X60-X84,Y87.0) | 12 | DNR | DNR | 10.0 - 35.7 |
| 10 Chronic liver disease and cirrhosis (K70,K73-K74) | 11 | DNR | DNR | 6.0 - 24.8 |
| Penobscot | | | | |
| 1 Cancer (C00-C97) | 379 | 250.8 | 178.5 | 160.2 - 196.8 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 347 | 229.7 | 161.3 | 144.2 - 178.4 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 120 | 79.4 | 71.1 | 57.7 - 84.6 |
| 4 Chronic lower respiratory disease (J40-J47) | 107 | 70.8 | 50.4 | 40.8 - 60.1 |
| 5 Cerebrovascular disease (I60-I69) | 72 | 47.7 | 33.6 | 26.2 - 42.4 |
| 6 Diabetes (E10-E14) | 61 | 40.4 | 28.4 | 21.6 - 36.7 |
| 7 Alzheimer's disease (G30) | 50 | 33.1 | 23.5 | 17.4 - 31.0 |
| 8 Suicide (X60-X84,Y87.0) | 40 | 26.5 | 23.6 | 16.4 - 32.8 |

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 3 of 5)

| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
|--|------------------|------------|-------------------|---------------|
| | | | Rate | 95% CI |
| Knox | | | | |
| 1 Cancer (C00-C97) | 134 | 336.9 | 179.9 | 148.9 - 210.9 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 112 | 281.6 | 154.5 | 125.5 - 183.4 |
| 3 Chronic lower respiratory disease (J40-J47) | 32 | 80.5 | 44.5 | 29.8 - 63.9 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 27 | 67.9 | 50.6 | 32.1 - 76.0 |
| 5 Alzheimer's disease (G30) | 21 | 52.8 | 28.7 | 17.8 - 43.9 |
| 6 Cerebrovascular disease (I60-I69) | 21 | 52.8 | 28.4 | 17.3 - 43.9 |
| 7 Diabetes (E10-E14) | 10 | DNR | DNR | 6.9 - 28.7 |
| Lincoln | | | | |
| 1 Heart disease (I00-I09,I11,I13,I20-I51) | 98 | 285.4 | 153.0 | 122.5 - 188.7 |
| 2 Cancer (C00-C97) | 96 | 279.5 | 138.4 | 111.3 - 170.1 |
| 3 Cerebrovascular disease (I60-I69) | 31 | 90.3 | 44.9 | 30.5 - 63.7 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 22 | 64.1 | 54.3 | 31.0 - 88.2 |
| 5 Alzheimer's disease (G30) | 20 | DNR | DNR | 18.0 - 45.5 |
| 6 Chronic lower respiratory disease (J40-J47) | 20 | DNR | DNR | 17.9 - 45.3 |
| 7 Diabetes (E10-E14) | 11 | DNR | DNR | 7.9 - 28.4 |
| Oxford | | | | |
| 1 Cancer (C00-C97) | 168 | 291.6 | 186.7 | 157.2 - 216.4 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 122 | 211.7 | 138.5 | 113.8 - 165.1 |
| 3 Chronic lower respiratory disease (J40-J47) | 59 | 102.4 | 64.4 | 48.8 - 83.4 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 51 | 88.5 | 82.3 | 59.8 - 110.5 |
| 5 Diabetes (E10-E14) | 30 | 52.1 | 32.6 | 21.9 - 46.9 |
| 6 Cerebrovascular disease (I60-I69) | 24 | 41.7 | 28.1 | 17.8 - 42.2 |
| 7 Alzheimer's disease (G30) | 21 | 36.4 | 24.3 | 15.1 - 37.2 |
| 8 Influenza and pneumonia (J10-J18) | 18 | DNR | DNR | 11.3 - 31.0 |
| 9 Suicide (X60-X84,Y87.0) | 13 | DNR | DNR | 11.2 - 37.8 |
| 10 Chronic liver disease and cirrhosis (K70,K73-K74) | 12 | DNR | DNR | 6.7 - 25.6 |
| Penobscot | | | | |
| 1 Cancer (C00-C97) | 377 | 249.5 | 177.0 | 158.8 - 195.2 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 351 | 232.3 | 163.2 | 146.0 - 180.4 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 120 | 79.4 | 71.1 | 57.7 - 84.6 |
| 4 Chronic lower respiratory disease (J40-J47) | 104 | 68.8 | 49.1 | 39.6 - 58.7 |
| 5 Cerebrovascular disease (I60-I69) | 73 | 48.3 | 34.0 | 26.6 - 42.8 |
| 6 Diabetes (E10-E14) | 63 | 41.7 | 29.5 | 22.6 - 37.9 |
| 7 Alzheimer's disease (G30) | 50 | 33.1 | 23.5 | 17.4 - 31.0 |
| 8 Suicide (X60-X84,Y87.0) | 41 | 27.1 | 24.3 | 17.0 - 33.6 |

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 4 of 5)

| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
|--|------------------|------------|-------------------|---------------|
| | | | Rate | 95% CI |
| Penobscot (continued) | | | | |
| 9 Influenza and pneumonia (J10-J18) | 38 | 25.1 | 17.2 | 12.1 - 23.8 |
| 10 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 28 | 18.5 | 13.6 | 9.0 - 19.8 |
| Piscataquis | | | | |
| 1 Cancer (C00-C97) | 63 | 375.0 | 195.6 | 148.9 - 252.3 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 58 | 345.2 | 179.8 | 135.5 - 234.0 |
| 3 Chronic lower respiratory disease (J40-J47) | 20 | DNR | DNR | 35.6 - 89.9 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 18 | DNR | DNR | 62.9 - 185.4 |
| Sagadahoc | | | | |
| 1 Cancer (C00-C97) | 96 | 269.4 | 169.0 | 136.0 - 207.4 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 86 | 241.3 | 157.0 | 125.6 - 193.9 |
| 3 Cerebrovascular disease (I60-I69) | 24 | 67.4 | 45.6 | 29.2 - 67.8 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 19 | DNR | DNR | 22.7 - 60.5 |
| 5 Chronic lower respiratory disease (J40-J47) | 19 | DNR | DNR | 20.2 - 52.4 |
| 6 Suicide (X60-X84,Y87.0) | 11 | DNR | DNR | 16.0 - 61.4 |
| 7 Alzheimer's disease (G30) | 11 | DNR | DNR | 9.9 - 35.7 |
| Somerset | | | | |
| 1 Cancer (C00-C97) | 138 | 272.8 | 180.9 | 149.6 - 212.2 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 138 | 272.8 | 188.1 | 156.0 - 220.1 |
| 3 Chronic lower respiratory disease (J40-J47) | 47 | 92.9 | 63.6 | 46.5 - 84.8 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 41 | 81.0 | 71.8 | 50.6 - 99.0 |
| 5 Cerebrovascular disease (I60-I69) | 18 | DNR | DNR | 15.2 - 40.5 |
| 6 Diabetes (E10-E14) | 15 | DNR | DNR | 10.5 - 32.4 |
| 7 Alzheimer's disease (G30) | 13 | DNR | DNR | 9.8 - 31.5 |
| 8 Suicide (X60-X84,Y87.0) | 13 | DNR | DNR | 11.7 - 39.7 |
| 9 Influenza and pneumonia (J10-J18) | 11 | DNR | DNR | 8.3 - 31.8 |
| Waldo | | | | |
| 1 Cancer (C00-C97) | 104 | 262.0 | 167.7 | 135.7 - 205.0 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 88 | 221.7 | 158.3 | 126.6 - 195.5 |
| 3 Chronic lower respiratory disease (J40-J47) | 20 | 50.4 | 32.3 | 19.7 - 49.8 |
| 4 Cerebrovascular disease (I60-I69) | 28 | 70.5 | 45.9 | 30.3 - 66.8 |
| 5 Unintentional injury (V01-X59,Y85-Y86) | 22 | 55.4 | 47.3 | 28.5 - 73.9 |
| 6 Diabetes (E10-E14) | 15 | DNR | DNR | 14.2 - 43.6 |
| 7 Alzheimer's disease (G30) | 14 | DNR | DNR | 13.3 - 40.9 |

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 4 of 5)

| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
|--|------------------|------------|-------------------|---------------|
| | | | Rate | 95% CI |
| Penobscot (continued) | | | | |
| 9 Influenza and pneumonia (J10-J18) | 37 | 24.5 | 17.4 | 12.1 - 24.2 |
| 10 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 28 | 18.5 | 13.6 | 9.0 - 19.8 |
| Piscataquis | | | | |
| 1 Cancer (C00-C97) | 57 | 339.3 | 178.1 | 133.4 - 232.9 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 54 | 321.4 | 169.1 | 125.9 - 222.3 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 17 | DNR | DNR | 56.4 - 172.9 |
| 4 Chronic lower respiratory disease (J40-J47) | 14 | DNR | DNR | 22.9 - 70.3 |
| Sagadahoc | | | | |
| 1 Cancer (C00-C97) | 96 | 269.4 | 169.0 | 136.0 - 207.4 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 86 | 241.3 | 157.0 | 125.6 - 193.9 |
| 3 Cerebrovascular disease (I60-I69) | 24 | 67.4 | 45.6 | 29.2 - 67.8 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 19 | DNR | DNR | 22.7 - 60.5 |
| 5 Chronic lower respiratory disease (J40-J47) | 19 | DNR | DNR | 20.2 - 52.4 |
| 6 Suicide (X60-X84,Y87.0) | 11 | DNR | DNR | 16.0 - 61.4 |
| 7 Alzheimer's disease (G30) | 11 | DNR | DNR | 9.9 - 35.7 |
| Somerset | | | | |
| 1 Heart disease (I00-I09,I11,I13,I20-I51) | 138 | 272.8 | 188.1 | 156.0 - 220.1 |
| 2 Cancer (C00-C97) | 138 | 272.8 | 180.9 | 149.6 - 212.2 |
| 3 Chronic lower respiratory disease (J40-J47) | 47 | 92.9 | 63.6 | 46.5 - 84.8 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 41 | 81.0 | 71.8 | 50.6 - 99.1 |
| 5 Cerebrovascular disease (I60-I69) | 18 | DNR | DNR | 15.2 - 40.5 |
| 6 Diabetes (E10-E14) | 15 | DNR | DNR | 10.5 - 32.4 |
| 7 Suicide (X60-X84,Y87.0) | 13 | DNR | DNR | 11.7 - 39.7 |
| 8 Alzheimer's disease (G30) | 13 | DNR | DNR | 9.8 - 31.5 |
| 9 Influenza and pneumonia (J10-J18) | 11 | DNR | DNR | 8.3 - 31.8 |
| Waldo | | | | |
| 1 Cancer (C00-C97) | 104 | 262.0 | 167.7 | 135.7 - 205.0 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 88 | 221.7 | 158.3 | 126.6 - 195.5 |
| 3 Cerebrovascular disease (I60-I69) | 28 | 70.5 | 45.9 | 30.3 - 66.8 |
| 4 Unintentional injury (V01-X59,Y85-Y86) | 22 | 55.4 | 47.3 | 28.5 - 73.9 |
| 5 Chronic lower respiratory disease (J40-J47) | 20 | DNR | DNR | 19.7 - 49.8 |
| 6 Diabetes (E10-E14) | 15 | DNR | DNR | 14.2 - 43.6 |
| 7 Alzheimer's disease (G30) | 14 | DNR | DNR | 13.3 - 40.9 |

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 5 of 5)

| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
|--|------------------|------------|-------------------|---------------|
| | | | Rate | 95% CI |
| Washington | | | | |
| 1 Heart disease (I00-I09,I11,I13,I20-I51) | 105 | 333.4 | 198.1 | 159.3 - 236.9 |
| 2 Cancer (C00-C97) | 86 | 273.1 | 154.4 | 123.0 - 191.4 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 33 | 104.8 | 93.1 | 61.4 - 135.5 |
| 4 Chronic lower respiratory disease (J40-J47) | 19 | 60.3 | 33.1 | 19.9 - 51.7 |
| 5 Cerebrovascular disease (I60-I69) | 22 | 69.9 | 43.7 | 27.1 - 66.9 |
| 6 Alzheimer's disease (G30) | 12 | DNR | DNR | 11.7 - 39.6 |
| 7 Suicide (*U03,X60-X84,Y87.0) | 12 | DNR | DNR | 17.4 - 66.8 |
| 8 Diabetes (E10-E14) | 12 | DNR | DNR | 10.3 - 34.7 |
| 9 Influenza and pneumonia (J10-J18) | 11 | DNR | DNR | 9.6 - 40.0 |
| 10 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 11 | DNR | DNR | 9.6 - 40.0 |
| York | | | | |
| 1 Cancer (C00-C97) | 494 | 239.5 | 157.9 | 143.6 - 172.1 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 412 | 198.8 | 131.7 | 118.8 - 144.5 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 181 | 87.8 | 77.7 | 65.7 - 89.7 |
| 4 Chronic lower respiratory disease (J40-J47) | 122 | 59.2 | 37.7 | 30.9 - 44.5 |
| 5 Alzheimer's disease (G30) | 92 | 44.6 | 28.4 | 22.9 - 34.8 |
| 6 Cerebrovascular disease (I60-I69) | 84 | 40.7 | 26.4 | 21.0 - 32.8 |
| 7 Diabetes (E10-E14) | 67 | 32.5 | 20.7 | 16.0 - 26.5 |
| 8 Influenza and pneumonia (J10-J18) | 50 | 24.2 | 16.2 | 11.9 - 21.4 |
| 9 Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 45 | 21.8 | 15.0 | 10.8 - 20.3 |
| 10 Suicide (X60-X84,Y87.0) | 37 | 17.9 | 15.5 | 10.6 - 21.9 |

Data Source: Maine CDC, Data Research, and Vital Statistics. March 2022.
 ICD-10 codes are grouped by ICD-10 113 Cause List.
 Crude rates are deaths per 100,000 population.
 Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.
 Cause of death categories with numerators less than 10 are suppressed by the WONDER system to protect privacy and are not reported on this table.
 Counties with fewer than 10 leading causes are due to small numbers. * ICD codes for terrorism are preceded with "**".

Table 6 (continued). Leading Causes of Death by County, Maine, 2018 (page 5 of 5)

| Cause of Death | Number of Deaths | Crude Rate | Age-Adjusted Rate | |
|---|------------------|------------|-------------------|---------------|
| | | | Rate | 95% CI |
| Washington | | | | |
| 1 Heart disease (I00-I09,I11,I13,I20-I51) | 101 | 320.7 | 191.4 | 155.0 - 233.7 |
| 2 Cancer (C00-C97) | 88 | 279.5 | 158.8 | 126.9 - 196.4 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 33 | 104.8 | 93.1 | 61.4 - 135.5 |
| 4 Chronic lower respiratory disease (J40-J47) | 22 | 69.9 | 38.3 | 24.0 - 57.9 |
| 5 Cerebrovascular disease (I60-I69) | 21 | 66.7 | 42.2 | 25.4 - 65.8 |
| 6 Alzheimer's disease (G30) | 12 | DNR | DNR | 11.7 - 39.6 |
| 7 Influenza and pneumonia (J10-J18) | 12 | DNR | DNR | 10.2 - 34.3 |
| 8 Suicide (*U03,X60-X84,Y87.0) | 11 | DNR | DNR | 14.7 - 61.1 |
| 9 Nephritis, nephrotic syndrome and nephrosis | 11 | DNR | DNR | 9.6 - 34.5 |
| 10 Diabetes (E10-E14) | 10 | DNR | DNR | 7.7 - 29.4 |
| York | | | | |
| 1 Cancer (C00-C97) | 494 | 239.5 | 157.9 | 143.6 - 172.1 |
| 2 Heart disease (I00-I09,I11,I13,I20-I51) | 412 | 198.8 | 131.7 | 118.8 - 144.5 |
| 3 Unintentional injury (V01-X59,Y85-Y86) | 181 | 87.8 | 77.7 | 65.7 - 89.7 |
| 4 Chronic lower respiratory disease (J40-J47) | 122 | 59.2 | 37.7 | 30.9 - 44.5 |
| 5 Alzheimer's disease (G30) | 92 | 44.6 | 28.4 | 22.9 - 34.8 |
| 6 Cerebrovascular disease (I60-I69) | 84 | 40.7 | 26.4 | 21.0 - 32.8 |
| 7 Diabetes (E10-E14) | 67 | 32.5 | 20.7 | 16.0 - 26.5 |
| 8 Influenza and pneumonia (J10-J18) | 50 | 24.2 | 16.2 | 11.9 - 21.4 |
| 9 Nephritis, nephrotic syndrome and nephrosis | 45 | 21.8 | 15.0 | 10.8 - 20.3 |
| 10 Suicide (X60-X84,Y87.0) | 37 | 17.9 | 15.5 | 10.6 - 21.9 |

Data Source: Maine CDC, Data Research, and Vital Statistics. March 2022.
 ICD-10 codes are grouped by ICD-10 113 Cause List.
 Crude rates are deaths per 100,000 population.
 Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 DNR: Data not reliable; death rates are suppressed as unreliable by the WONDER system when the numerator is 20 or less.
 Cause of death categories with numerators less than 10 are suppressed by the WONDER system to protect privacy and are not reported on this table.
 * ICD codes for terrorism are preceded with "**".

Table 12. All Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|---------------|-------------------|----------------------|
| | | | Rate | 95% CI |
| Maine total | 14,302 | 1072.9 | 760.8 | 755.2 - 766.4 |
| County | | | | |
| Androscoggin | 1,140 | 1060.6 | 837.4 | 815.9 - 858.8 |
| Aroostook | 875 | 1283.0 | 793.8 | 769.6 - 818.0 |
| Cumberland | 2,666 | 915.2 | 687.3 | 675.7 - 698.9 |
| Franklin | 346 | 1153.0 | 806.2 | 768.1 - 844.3 |
| Hancock | 612 | 1122.9 | 714.8 | 688.9 - 740.8 |
| Kennebec | 1,337 | 1103.5 | 808.0 | 788.6 - 827.3 |
| Knox | 492 | 1237.2 | 731.3 | 701.5 - 761.1 |
| Lincoln | 426 | 1244.8 | 711.5 | 679.5 - 743.5 |
| Oxford | 653 | 1138.3 | 777.3 | 750.5 - 804.1 |
| Penobscot | 1,632 | 1072.4 | 808.5 | 791.0 - 826.0 |
| Piscataquis | 256 | 1519.5 | 919.4 | 866.7 - 972.1 |
| Sagadahoc | 371 | 1052.0 | 717.6 | 684.9 - 750.4 |
| Somerset | 609 | 1197.2 | 868.7 | 837.7 - 899.6 |
| Waldo | 406 | 1030.1 | 744.6 | 712.0 - 777.2 |
| Washington | 433 | 1370.7 | 896.7 | 857.8 - 935.6 |
| York | 2,046 | 1008.4 | 721.2 | 707.1 - 735.2 |

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019.
 All Deaths: Deaths in which the underlying cause of death was coded as ICD-10 any listed.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 12. All Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|---------------|-------------------|----------------------|
| | | | Rate | 95% CI |
| Maine total | 14,304 | 1073.0 | 760.9 | 755.3 - 766.5 |
| County | | | | |
| Androscoggin | 1,140 | 1060.8 | 837.5 | 816.0 - 858.9 |
| Aroostook | 875 | 1282.7 | 793.9 | 769.7 - 818.1 |
| Cumberland | 2,666 | 915.4 | 687.5 | 675.9 - 699.1 |
| Franklin | 346 | 1152.5 | 805.9 | 767.8 - 843.9 |
| Hancock | 610 | 1119.2 | 712.4 | 686.5 - 738.2 |
| Kennebec | 1,337 | 1103.9 | 808.1 | 788.8 - 827.5 |
| Knox | 492 | 1237.0 | 731.2 | 701.4 - 761.0 |
| Lincoln | 426 | 1244.8 | 711.9 | 679.8 - 743.9 |
| Oxford | 666 | 1162.3 | 793.0 | 765.9 - 820.0 |
| Penobscot | 1,630 | 1071.3 | 808.5 | 791.0 - 826.0 |
| Piscataquis | 243 | 1437.4 | 871.3 | 819.9 - 922.7 |
| Sagadahoc | 371 | 1052.0 | 717.5 | 684.8 - 750.2 |
| Somerset | 609 | 1197.4 | 869.2 | 838.2 - 900.2 |
| Waldo | 408 | 1034.7 | 748.3 | 715.6 - 781.0 |
| Washington | 434 | 1375.3 | 894.8 | 856.2 - 933.5 |
| York | 2,047 | 1008.9 | 721.5 | 707.4 - 735.6 |

Data Source: Maine CDC, Data, Research and Vital Statistics. March 2022.
 All Deaths: Deaths in which the underlying cause of death was coded as ICD-10 any listed.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 17. Cancer Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|--------------|-------------------|----------------------|
| | | | Rate | 95% CI |
| Maine total | 3,304 | 247.9 | 169.6 | 167.0 - 172.2 |
| County | | | | |
| Androscoggin | 250 | 233.0 | 178.4 | 168.4 - 188.4 |
| Aroostook | 186 | 272.8 | 166.8 | 155.6 - 178.0 |
| Cumberland | 599 | 205.8 | 153.5 | 147.9 - 159.1 |
| Franklin | 81 | 270.9 | 181.0 | 163.0 - 199.1 |
| Hancock | 142 | 259.6 | 157.8 | 145.8 - 169.8 |
| Kennebec | 307 | 253.6 | 176.5 | 167.5 - 185.5 |
| Knox | 120 | 300.9 | 175.0 | 160.4 - 189.5 |
| Lincoln | 100 | 292.5 | 156.4 | 142.0 - 170.8 |
| Oxford | 157 | 274.2 | 179.4 | 166.5 - 192.3 |
| Penobscot | 356 | 234.2 | 171.0 | 162.9 - 179.1 |
| Piscataquis | 66 | 393.5 | 215.6 | 191.8 - 239.5 |
| Sagadahoc | 93 | 264.6 | 170.2 | 154.4 - 185.9 |
| Somerset | 147 | 289.4 | 196.7 | 182.1 - 211.3 |
| Waldo | 97 | 245.6 | 165.5 | 150.3 - 180.8 |
| Washington | 103 | 325.4 | 197.6 | 179.8 - 215.3 |
| York | 498 | 245.3 | 169.5 | 162.7 - 176.3 |

Data Source: Maine CDC, Data Research and Vital statistics. March 2020
 Cancer Deaths: Deaths with cancer as the underlying cause of death.
 Cancer: ICD-10 codes: C00-C97
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 17. Cancer Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|--------------|-------------------|----------------------|
| | | | Rate | 95% CI |
| Maine total | 3,304 | 247.8 | 169.6 | 166.9 - 172.2 |
| County | | | | |
| Androscoggin | 250 | 233.0 | 178.4 | 168.4 - 188.4 |
| Aroostook | 186 | 272.7 | 166.7 | 155.6 - 177.9 |
| Cumberland | 599 | 205.8 | 153.5 | 147.9 - 159.1 |
| Franklin | 81 | 270.8 | 181.0 | 163.0 - 199.1 |
| Hancock | 142 | 260.0 | 158.0 | 146.0 - 170.0 |
| Kennebec | 307 | 253.6 | 176.5 | 167.5 - 185.5 |
| Knox | 120 | 300.8 | 175.0 | 160.4 - 189.5 |
| Lincoln | 100 | 292.5 | 156.4 | 142.1 - 170.8 |
| Oxford | 161 | 281.2 | 183.7 | 170.7 - 196.7 |
| Penobscot | 356 | 233.7 | 170.7 | 162.6 - 178.7 |
| Piscataquis | 62 | 369.7 | 203.1 | 179.9 - 226.3 |
| Sagadahoc | 93 | 264.6 | 170.2 | 154.4 - 185.9 |
| Somerset | 147 | 289.3 | 196.7 | 182.1 - 211.3 |
| Waldo | 97 | 245.1 | 165.3 | 150.0 - 180.5 |
| Washington | 104 | 328.0 | 198.8 | 181.0 - 216.5 |
| York | 498 | 245.3 | 169.5 | 162.7 - 176.3 |

Data Source: Maine CDC, Data Research and Vital statistics. March 2022.
 Cancer Deaths: Deaths with cancer as the underlying cause of death.
 Cancer: ICD-10 codes C00-C97
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 22. Heart Disease Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|--------------|-------------------|----------------------|
| | | | Rate | 95% CI |
| Maine total | 2,889 | 216.8 | 148.5 | 146.1 - 151.0 |
| County | | | | |
| Androscoggin | 231 | 215.0 | 166.0 | 156.4 - 175.6 |
| Aroostook | 187 | 274.8 | 160.0 | 149.5 - 170.5 |
| Cumberland | 501 | 172.0 | 124.8 | 119.9 - 129.8 |
| Franklin | 68 | 225.7 | 152.8 | 136.3 - 169.2 |
| Hancock | 126 | 231.0 | 142.8 | 131.4 - 154.2 |
| Kennebec | 270 | 222.5 | 158.7 | 150.2 - 167.3 |
| Knox | 112 | 280.7 | 153.3 | 140.4 - 166.2 |
| Lincoln | 92 | 270.3 | 147.2 | 133.2 - 161.2 |
| Oxford | 119 | 206.9 | 138.5 | 127.2 - 149.8 |
| Penobscot | 334 | 219.7 | 161.1 | 153.3 - 168.9 |
| Piscataquis | 56 | 333.1 | 190.4 | 167.3 - 213.5 |
| Sagadahoc | 80 | 225.5 | 147.9 | 133.3 - 162.5 |
| Somerset | 140 | 274.4 | 195.1 | 180.5 - 209.7 |
| Waldo | 94 | 239.5 | 169.9 | 154.4 - 185.4 |
| Washington | 91 | 288.7 | 179.3 | 162.5 - 196.2 |
| York | 388 | 191.4 | 131.6 | 125.7 - 137.5 |

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019.
 Heart Disease Deaths: Deaths with heart disease as the underlying cause of death.
 Heart Disease: ICD-10 codes I00-I09, I11, I13, I20-I51.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 22. Heart Disease Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|--------------|-------------------|----------------------|
| | | | Rate | 95% CI |
| Maine total | 2,890 | 216.8 | 148.5 | 146.1 - 151.0 |
| County | | | | |
| Androscoggin | 231 | 214.9 | 165.9 | 156.3 - 175.6 |
| Aroostook | 187 | 274.8 | 160.0 | 149.5 - 170.4 |
| Cumberland | 501 | 172.0 | 124.8 | 119.9 - 129.8 |
| Franklin | 68 | 225.6 | 152.7 | 136.3 - 169.1 |
| Hancock | 126 | 230.7 | 142.6 | 131.2 - 154.0 |
| Kennebec | 270 | 222.7 | 158.8 | 150.3 - 167.4 |
| Knox | 112 | 280.7 | 153.3 | 140.4 - 166.2 |
| Lincoln | 92 | 270.3 | 147.2 | 133.3 - 161.2 |
| Oxford | 121 | 210.7 | 141.1 | 129.7 - 152.5 |
| Penobscot | 334 | 219.3 | 161.0 | 153.2 - 168.8 |
| Piscataquis | 54 | 319.9 | 182.5 | 160.0 - 205.0 |
| Sagadahoc | 80 | 225.5 | 147.9 | 133.3 - 162.5 |
| Somerset | 140 | 274.4 | 195.1 | 180.4 - 209.7 |
| Waldo | 94 | 239.5 | 169.8 | 154.3 - 185.3 |
| Washington | 92 | 290.0 | 180.1 | 163.2 - 197.0 |
| York | 388 | 191.4 | 131.6 | 125.7 - 137.5 |

Data Source: Maine CDC, Data, Research and Vital Statistics. March 2022.
 Heart Disease Deaths: Deaths with heart disease as the underlying cause of death.
 Heart Disease: ICD-10 codes I00-I09, I11, I13, I20-I51.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 27. Chronic Lower Respiratory Disease Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|-------------|-------------------|--------------------|
| | | | Rate | 95% CI |
| Maine total | 951 | 71.4 | 48.5 | 47.1 - 49.9 |
| County | | | | |
| Androscoggin | 80 | 74.1 | 57.4 | 51.7 - 63.1 |
| Aroostook | 69 | 101.8 | 59.2 | 52.9 - 65.6 |
| Cumberland | 154 | 52.8 | 38.4 | 35.6 - 41.1 |
| Franklin | 25 | 84.5 | 55.1 | 45.4 - 64.8 |
| Hancock | 37 | 68.2 | 41.8 | 35.6 - 47.9 |
| Kennebec | 89 | 73.8 | 51.7 | 46.9 - 56.6 |
| Knox | 30 | 76.5 | 44.5 | 37.1 - 51.9 |
| Lincoln | 24 | 69.0 | 35.5 | 29.0 - 42.0 |
| Oxford | 58 | 101.5 | 65.9 | 58.2 - 73.6 |
| Penobscot | 115 | 75.7 | 56.1 | 51.5 - 60.8 |
| Piscataquis | 17 | 103.1 | 56.3 | 44.9 - 69.8 |
| Sagadahoc | 22 | 62.9 | 40.0 | 32.5 - 47.6 |
| Somerset | 50 | 99.1 | 69.0 | 60.3 - 77.6 |
| Waldo | 25 | 63.4 | 42.0 | 34.5 - 49.5 |
| Washington | 27 | 86.7 | 53.3 | 44.1 - 62.5 |
| York | 126 | 62.4 | 42.1 | 38.9 - 45.5 |

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019
 Chronic Lower Respiratory Disease (CLRD) Deaths: Deaths with CLRD as the underlying cause of death.
 Chronic Lower Respiratory Disease: ICD-10 codes J40-J47.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 27. Chronic Lower Respiratory Disease Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|-------------|-------------------|--------------------|
| | | | Rate | 95% CI |
| Maine total | 951 | 71.4 | 48.5 | 47.1 - 49.9 |
| County | | | | |
| Androscoggin | 80 | 74.1 | 57.4 | 51.7 - 63.1 |
| Aroostook | 69 | 101.8 | 59.2 | 52.9 - 65.6 |
| Cumberland | 154 | 52.8 | 38.4 | 35.6 - 41.1 |
| Franklin | 25 | 84.5 | 55.1 | 45.4 - 64.8 |
| Hancock | 37 | 67.5 | 41.3 | 35.2 - 47.5 |
| Kennebec | 89 | 73.8 | 51.7 | 46.9 - 56.6 |
| Knox | 30 | 76.5 | 44.5 | 37.1 - 51.9 |
| Lincoln | 24 | 69.0 | 35.5 | 29.0 - 42.0 |
| Oxford | 60 | 104.0 | 67.4 | 59.7 - 75.2 |
| Penobscot | 114 | 75.2 | 55.8 | 51.1 - 60.4 |
| Piscataquis | 16 | 94.8 | 51.9 | 41.0 - 64.9 |
| Sagadahoc | 22 | 62.9 | 40.0 | 32.5 - 47.6 |
| Somerset | 50 | 99.1 | 69.0 | 60.3 - 77.6 |
| Waldo | 25 | 64.4 | 42.7 | 35.1 - 50.2 |
| Washington | 28 | 89.3 | 54.8 | 45.5 - 64.1 |
| York | 127 | 62.4 | 42.2 | 38.9 - 45.5 |

Data Source: Maine CDC, Data, Research and Vital Statistics. March 2022.
 Chronic Lower Respiratory Disease (CLRD) Deaths: Deaths with CLRD as the underlying cause of death.
 Chronic Lower Respiratory Disease: ICD-10 codes J40-J47.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 66

Table 32. Unintentional Injury Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|--------------|
| | | | Rate | 95% CI |
| Maine total | 874 | 65.6 | 58.8 | 57.0 - 60.7 |
| County | | | | |
| Androscoggin | 66 | 61.0 | 56.9 | 50.6 - 63.3 |
| Aroostook | 38 | 55.7 | 47.3 | 40.0 - 54.7 |
| Cumberland | 173 | 59.5 | 52.8 | 49.2 - 56.5 |
| Franklin | 18 | 61.2 | 55.4 | 43.9 - 68.9 |
| Hancock | 37 | 68.2 | 59.0 | 49.8 - 68.2 |
| Kennebec | 82 | 67.7 | 62.2 | 55.9 - 68.5 |
| Knox | 30 | 76.0 | 60.3 | 55.9 - 68.5 |
| Lincoln | 23 | 68.4 | 58.4 | 47.0 - 71.7 |
| Oxford | 34 | 58.6 | 51.9 | 43.5 - 60.4 |
| Penobscot | 99 | 64.8 | 60.0 | 54.5 - 65.6 |
| Piscataquis | 15 | 91.3 | 82.5 | 63.3 - 105.8 |
| Sagadahoc | 18 | 50.4 | 41.1 | 32.5 - 51.2 |
| Somerset | 37 | 71.9 | 66.7 | 56.5 - 76.9 |
| Waldo | 25 | 64.4 | 59.4 | 48.4 - 70.3 |
| Washington | 32 | 101.3 | 98.6 | 82.0 - 115.1 |
| York | 146 | 71.8 | 65.6 | 60.6 - 70.6 |

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019. Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. Unintentional Injury Deaths: Deaths with unintentional injury as the underlying cause of death. Unintentional Injury: ICD-10 codes V01-X59,Y85-Y86. Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population. 95% CI: 95% confidence interval of the rate. Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 32. Unintentional Injury Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|--------------|
| | | | Rate | 95% CI |
| Maine total | 874 | 65.5 | 58.7 | 56.9 - 60.6 |
| County | | | | |
| Androscoggin | 66 | 61.0 | 56.9 | 50.6 - 63.3 |
| Aroostook | 38 | 55.7 | 47.3 | 40.0 - 54.7 |
| Cumberland | 173 | 59.5 | 52.8 | 49.2 - 56.5 |
| Franklin | 18 | 61.2 | 55.4 | 43.9 - 68.9 |
| Hancock | 37 | 67.8 | 58.8 | 49.6 - 68.0 |
| Kennebec | 82 | 67.7 | 62.2 | 55.8 - 68.5 |
| Knox | 30 | 76.0 | 60.3 | 49.7 - 70.8 |
| Lincoln | 23 | 68.4 | 58.4 | 47.0 - 71.8 |
| Oxford | 35 | 60.3 | 53.3 | 44.8 - 61.9 |
| Penobscot | 100 | 65.8 | 60.8 | 55.3 - 66.4 |
| Piscataquis | 14 | 85.3 | 77.6 | 58.9 - 100.3 |
| Sagadahoc | 18 | 50.4 | 41.0 | 32.5 - 51.1 |
| Somerset | 37 | 71.9 | 66.7 | 56.5 - 76.9 |
| Waldo | 26 | 65.0 | 59.7 | 48.8 - 70.7 |
| Washington | 30 | 96.2 | 94.4 | 78.1 - 110.6 |
| York | 146 | 71.9 | 65.6 | 60.6 - 70.6 |

Data Source: Maine CDC, Data, Research and Vital Statistics. March 2022. Unintentional Injury Deaths: Deaths with unintentional injury as the underlying cause of death. Unintentional Injury: ICD-10 codes V01-X59,Y85-Y86. Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population. 95% CI: 95% confidence interval of the rate. Subgroup counts might not sum to Maine total due to missing data or rounding.

Page 70

Table 37. Cerebrovascular Disease Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|-------------|
| | | | Rate | 95% CI |
| Maine total | 664 | 49.8 | 34.2 | 33.0 - 35.4 |
| County | | | | |
| Androscoggin | 54 | 49.9 | 38.7 | 34.0 - 43.4 |
| Aroostook | 41 | 59.8 | 34.9 | 30.0 - 39.7 |
| Cumberland | 127 | 43.7 | 31.8 | 29.3 - 34.3 |
| Franklin | 18 | 60.6 | 39.9 | 32.1 - 49.1 |
| Hancock | 31 | 57.6 | 34.3 | 28.8 - 39.8 |
| Kennebec | 69 | 57.3 | 40.9 | 36.5 - 45.2 |
| Knox | 22 | 55.8 | 30.4 | 24.6 - 36.3 |
| Lincoln | 28 | 82.5 | 42.3 | 35.2 - 49.5 |
| Oxford | 24 | 42.2 | 28.1 | 23.0 - 33.1 |
| Penobscot | 71 | 46.9 | 34.5 | 30.9 - 38.1 |
| Piscataquis | 12 | 69.9 | 40.9 | 30.7 - 53.6 |
| Sagadahoc | 24 | 68.0 | 45.7 | 37.4 - 54.0 |
| Somerset | 24 | 46.4 | 33.8 | 27.7 - 40.0 |
| Waldo | 21 | 52.3 | 37.4 | 30.4 - 45.6 |
| Washington | 16 | 51.3 | 32.3 | 25.5 - 40.4 |
| York | 81 | 39.9 | 27.7 | 25.0 - 30.4 |

Data Source: Maine CDC, Data, Research and Vital Statistics. December 2019. Cerebrovascular Disease Deaths: Deaths with cerebrovascular disease as the underlying cause of death. Cerebrovascular Disease: ICD-10 codes I60-I69. Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population. 95% CI: 95% confidence interval of the rate. Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 37. Cerebrovascular Disease Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|-------------|
| | | | Rate | 95% CI |
| Maine total | 664 | 49.8 | 34.2 | 33.0 - 35.3 |
| County | | | | |
| Androscoggin | 54 | 49.9 | 38.7 | 34.0 - 43.4 |
| Aroostook | 41 | 59.8 | 34.9 | 30.0 - 39.7 |
| Cumberland | 127 | 43.7 | 31.8 | 29.3 - 34.3 |
| Franklin | 18 | 60.6 | 39.9 | 32.1 - 49.0 |
| Hancock | 32 | 57.9 | 34.5 | 29.0 - 40.0 |
| Kennebec | 69 | 57.3 | 40.9 | 36.5 - 45.2 |
| Knox | 22 | 55.8 | 30.4 | 24.6 - 36.3 |
| Lincoln | 28 | 82.5 | 42.4 | 35.2 - 49.5 |
| Oxford | 25 | 44.0 | 29.5 | 24.2 - 34.7 |
| Penobscot | 71 | 46.9 | 34.4 | 30.8 - 38.0 |
| Piscataquis | 11 | 64.0 | 36.3 | 27.0 - 47.9 |
| Sagadahoc | 24 | 68 | 45.7 | 37.5 - 54.0 |
| Somerset | 24 | 46.4 | 33.8 | 27.7 - 40.0 |
| Waldo | 20 | 51.8 | 37.1 | 30.1 - 45.2 |
| Washington | 16 | 51.3 | 32.3 | 25.4 - 40.4 |
| York | 81 | 39.9 | 27.7 | 25.0 - 30.4 |

Data Source: Maine CDC, Data, Research and Vital Statistics. March 2022. Cerebrovascular Disease Deaths: Deaths with cerebrovascular disease as the underlying cause of death. Cerebrovascular Disease: ICD-10 codes I60-I69. Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population. 95% CI: 95% confidence interval of the rate. Subgroup counts might not sum to Maine total due to missing data or rounding.

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|-------------|
| | | | Rate | 95% CI |
| | | | | |
| Maine total | 286 | 21.5 | 14.7 | 13.9 - 15.5 |
| County | | | | |
| Androscoggin | 25 | 22.7 | 17.3 | 14.2 - 20.4 |
| Aroostook | 20 | 29.3 | 17.0 | 13.8 - 20.8 |
| Cumberland | 54 | 18.7 | 13.4 | 11.8 - 15.1 |
| Franklin | 8 | 25.3 | 19.1 | 13.3 - 26.6 |
| Hancock | 13 | 23.1 | 14.0 | 10.6 - 18.1 |
| Kennebec | 24 | 19.5 | 14.1 | 11.5 - 16.6 |
| Knox | 7 | 18.6 | 10.6 | 7.2 - 14.9 |
| Lincoln | 8 | 22.2 | 11.6 | 8.1 - 16.1 |
| Oxford | 12 | 21.3 | 14.1 | 10.8 - 18.2 |
| Penobscot | 37 | 24.6 | 17.9 | 15.3 - 20.4 |
| Piscataquis | 4 | 23.7 | 14.1 | 8.2 - 22.6 |
| Sagadahoc | 8 | 22.7 | 14.6 | 10.5 - 19.9 |
| Somerset | 9 | 18.5 | 13.8 | 10.0 - 18.4 |
| Waldo | 8 | 20.3 | 14.9 | 10.6 - 20.3 |
| Washington | 12 | 36.7 | 21.7 | 16.3 - 28.2 |
| York | 38 | 18.6 | 12.9 | 11.1 - 14.8 |

Data Source: Maine CDC, Data, Research, and Vital Statistics. March 2020.
 Influenza and Pneumonia Deaths: Deaths with influenza or pneumonia as the underlying cause of death.
 Influenza and Pneumonia: ICD-10 codes J10-J18.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|-------------|
| | | | Rate | 95% CI |
| | | | | |
| Maine total | 285 | 21.4 | 14.6 | 13.9 - 15.4 |
| County | | | | |
| Androscoggin | 24 | 22.5 | 17.1 | 14.0 - 20.2 |
| Aroostook | 20 | 29.3 | 17.0 | 13.8 - 20.8 |
| Cumberland | 54 | 18.7 | 13.4 | 11.8 - 15.1 |
| Franklin | 8 | 25.3 | 19.1 | 13.3 - 26.6 |
| Hancock | 12 | 22.7 | 13.5 | 10.3 - 17.5 |
| Kennebec | 23 | 19.1 | 13.7 | 11.2 - 16.2 |
| Knox | 7 | 18.6 | 10.6 | 7.2 - 14.9 |
| Lincoln | 8 | 22.2 | 11.6 | 8.1 - 16.1 |
| Oxford | 12 | 21.3 | 14.1 | 10.8 - 18.2 |
| Penobscot | 37 | 24.4 | 17.9 | 15.3 - 20.5 |
| Piscataquis | 4 | 23.7 | 14.1 | 8.2 - 22.6 |
| Sagadahoc | 8 | 22.7 | 14.7 | 10.5 - 20.0 |
| Somerset | 9 | 18.1 | 13.5 | 9.8 - 18.2 |
| Waldo | 8 | 20.3 | 14.9 | 10.6 - 20.3 |
| Washington | 12 | 37.4 | 21.4 | 16.3 - 27.7 |
| York | 38 | 18.6 | 12.9 | 11.1 - 14.8 |

Data Source: Maine CDC, Data, Research, and Vital Statistics. March 2022.
 Influenza and Pneumonia Deaths: Deaths with influenza or pneumonia as the underlying cause of death.
 Influenza and Pneumonia: ICD-10 codes J10-J18.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 57. Suicide Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|-------------|
| | | | Rate | 95% CI |
| Maine total | 244 | 18.3 | 16.9 | 15.9 - 17.9 |
| County | | | | |
| Androscoggin | 20 | 18.8 | 17.8 | 14.4 - 21.7 |
| Aroostook | 14 | 19.9 | 18.7 | 14.2 - 24.2 |
| Cumberland | 38 | 13.2 | 12.3 | 10.5 - 14.1 |
| Franklin | 6 | 20.6 | 19.0 | 12.5 - 27.6 |
| Hancock | 10 | 18.7 | 15.8 | 11.4 - 21.2 |
| Kennebec | 25 | 20.5 | 19.3 | 15.8 - 22.9 |
| Knox | 8 | 19.6 | 17.9 | 12.3 - 25.1 |
| Lincoln | 7 | 20.5 | 20.2 | 13.6 - 28.8 |
| Oxford | 10 | 17.8 | 15.2 | 11.0 - 20.4 |
| Penobscot | 27 | 18.0 | 16.9 | 13.9 - 19.8 |
| Piscataquis | 4 | 23.7 | 23.2 | 13.5 - 37.1 |
| Sagadahoc | 9 | 24.4 | 23.2 | 16.4 - 31.8 |
| Somerset | 12 | 23.2 | 20.9 | 15.6 - 27.4 |
| Waldo | 9 | 22.3 | 19.6 | 13.9 - 26.8 |
| Washington | 9 | 27.2 | 26.0 | 18.2 - 36.0 |
| York | 36 | 17.9 | 16.8 | 14.2 - 19.4 |

Data Source: Maine CDC. Data, Research, Vital Statistics. December 2019
 Suicide Deaths: Deaths with suicide as the underlying cause of death.
 Suicide: ICD-10 codes X60-X84, Y87.0 .
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 57. Suicide Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|-------------|
| | | | Rate | 95% CI |
| Maine total | 244 | 18.3 | 16.9 | 15.9 - 17.9 |
| County | | | | |
| Androscoggin | 20 | 18.8 | 17.8 | 14.4 - 21.7 |
| Aroostook | 14 | 19.9 | 18.7 | 14.2 - 24.2 |
| Cumberland | 38 | 13.2 | 12.3 | 10.5 - 14.1 |
| Franklin | 6 | 20.6 | 19.0 | 12.5 - 27.6 |
| Hancock | 10 | 18.3 | 15.3 | 11.0 - 20.7 |
| Kennebec | 25 | 20.5 | 19.3 | 15.8 - 22.9 |
| Knox | 8 | 19.6 | 17.9 | 12.3 - 25.1 |
| Lincoln | 7 | 20.5 | 20.2 | 13.6 - 28.8 |
| Oxford | 11 | 18.8 | 15.8 | 11.6 - 21.1 |
| Penobscot | 28 | 18.4 | 17.2 | 14.2 - 20.2 |
| Piscataquis | 3 | 20.1 | 21.0 | 11.5 - 35.2 |
| Sagadahoc | 9 | 24.4 | 23.2 | 16.4 - 31.8 |
| Somerset | 12 | 23.2 | 20.9 | 15.6 - 27.4 |
| Waldo | 9 | 22.8 | 20.3 | 14.4 - 27.7 |
| Washington | 8 | 25.3 | 24.5 | 17.0 - 34.3 |
| York | 36 | 17.8 | 16.7 | 14.1 - 19.3 |

Data Source: Maine CDC. Data, Research, Vital Statistics. December March 2022.
 Suicide Deaths: Deaths with suicide as the underlying cause of death.
 Suicide: ICD-10 codes X60-X84, Y87.0 .
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 62. Nephritis, Nephrotic Syndrome, & Nephrosis Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|-------------|
| | | | Rate | 95% CI |
| Maine total | 238 | 17.9 | 12.2 | 11.5 - 12.9 |
| County | | | | |
| Androscoggin | 12 | 11.5 | 8.7 | 6.6 - 11.2 |
| Aroostook | 18 | 26.4 | 15.3 | 12.3 - 18.9 |
| Cumberland | 48 | 16.5 | 12.1 | 10.5 - 13.6 |
| Franklin | 7 | 22.0 | 14.9 | 10.3 - 21.0 |
| Hancock | 12 | 21.6 | 12.8 | 9.7 - 16.5 |
| Kennebec | 19 | 15.8 | 11.1 | 9.0 - 13.6 |
| Knox | 7 | 18.1 | 9.8 | 6.8 - 13.6 |
| Lincoln | 5 | 14.0 | 7.6 | 4.7 - 11.6 |
| Oxford | 12 | 20.6 | 12.9 | 9.8 - 16.7 |
| Penobscot | 34 | 22.1 | 16.2 | 13.7 - 18.6 |
| Piscataquis | 3 | 15.4 | 8.4 | 4.3 - 14.7 |
| Sagadahoc | 7 | 21.0 | 13.7 | 9.7 - 18.9 |
| Somerset | 6 | 11.8 | 8.1 | 5.4 - 11.6 |
| Waldo | 5 | 13.7 | 10.6 | 6.9 - 15.7 |
| Washington | 7 | 22.2 | 14.1 | 9.6 - 19.9 |
| York | 36 | 17.8 | 12.3 | 10.5 - 14.2 |

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics.
 Nephritis, Nephrotic Syndrome and Nephrosis Deaths: Deaths with nephritis, nephrotic syndrome or nephrosis as the underlying cause of death.
 Nephritis, Nephrotic Syndrome and Nephrosis: ICD-10 codes N00-N07, N17-N19, N25-N27.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.

Table 62. Nephritis, Nephrotic Syndrome, & Nephrosis Deaths by County, Maine, 2014 - 2018

| | Average Annual Number | Crude Rate | Age-Adjusted Rate | |
|--------------------|-----------------------|------------|-------------------|-------------|
| | | | Rate | 95% CI |
| Maine total | 238 | 17.9 | 12.2 | 11.5 - 12.9 |
| County | | | | |
| Androscoggin | 12 | 11.5 | 8.7 | 6.6 - 11.2 |
| Aroostook | 18 | 26.4 | 15.3 | 12.3 - 18.9 |
| Cumberland | 48 | 16.5 | 12.1 | 10.5 - 13.6 |
| Franklin | 7 | 22.0 | 14.9 | 10.3 - 20.9 |
| Hancock | 12 | 21.6 | 12.8 | 9.7 - 16.5 |
| Kennebec | 19 | 15.8 | 11.1 | 9.0 - 13.6 |
| Knox | 7 | 18.1 | 9.8 | 6.8 - 13.6 |
| Lincoln | 5 | 14.0 | 7.6 | 4.7 - 11.6 |
| Oxford | 12 | 21.3 | 13.3 | 10.2 - 17.2 |
| Penobscot | 34 | 22.2 | 16.4 | 13.9 - 18.9 |
| Piscataquis | 2 | 13.0 | 7.2 | 3.5 - 13.3 |
| Sagadahoc | 7 | 21.0 | 13.8 | 9.7 - 19.0 |
| Somerset | 6 | 11.8 | 8.1 | 5.4 - 11.6 |
| Waldo | 5 | 13.7 | 10.6 | 6.9 - 15.7 |
| Washington | 7 | 21.5 | 13.1 | 9.0 - 18.4 |
| York | 36 | 17.8 | 12.3 | 10.5 - 14.2 |

Data Source: Maine Center for Disease Control and Prevention. Data, Research and Vital Statistics. March 2022.
 Nephritis, Nephrotic Syndrome and Nephrosis Deaths: Deaths with nephritis, nephrotic syndrome or nephrosis as the underlying cause of death.
 Nephritis, Nephrotic Syndrome and Nephrosis: ICD-10 codes N00-N07, N17-N19, N25-N27.
 Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population.
 95% CI: 95% confidence interval of the rate.
 Subgroup counts might not sum to Maine total due to missing data or rounding.